

Lecture: 10am, Wednesday 23rd November

MAS2906: Computational Probability and Statistics with R

Newcastle University
`lee.fawcett@ncl.ac.uk`

Semester 1, 2022/23

Assessments

- Numbas test 1 – will make a short video going through the questions very soon, with feedback on your work

■ Mini Project

- Now available in Canvas
- Due 4pm, Friday 9th December
- You can work in pairs if you like!
- Standard submission via Canvas; no Numbas; any code should be included in your solutions document
- Will give you help with this in the practical sessions this Friday – do come along!

Assessments

- Numbas test 1 – will make a short video going through the questions very soon, with feedback on your work
- Mini Project
 - Now available in Canvas
 - Due 4pm, Friday 9th December
 - You can work in pairs if you like!
 - Standard submission via Canvas; no Numbas; any code should be included in your solutions document
 - Will give you help with this in the practical sessions this Friday – do come along!

Chapter 5

Transformations

5.1 Transforming a normal random variable

- Consider how we might model a **continuous positive** random quantity e.g. blood pressure or height
- The Normal distribution is not suitable!


One way to obtain a positive random variable is to define

$$Y = \exp(X), \quad \text{where } X \sim N(\mu, \sigma^2).$$

Then Y is a random variable which is a **transformation** of X .

5.1 Simulation to explore transformation

```
1 y = exp(rnorm(250, mean = 1.5, sd = 0.3))  
2 hist(y, xlab = 'Y', ylab = 'frequency')
```



5.1 Calculating PDFs

- This looks like a useful distribution – what is its PDF?
- $Y = \exp(X)$ so $X = \log(Y)$
- So $f_Y(y) = f_X(\log(y))$ is a good guess, but is **NOT** the right answer!

Example

If $X \sim U(0, 1)$, what is the PDF of $Y = X^3$?

Example 5.1: Solution

$$\begin{aligned}F_Y(y) &= \Pr(Y \leq y) \\&= \Pr(X^3 \leq y) \\&= \Pr(X \leq y^{1/3}) \\&= F_X(y^{1/3}).\end{aligned}$$

Therefore

$$\begin{aligned}f_Y(y) &= \frac{d}{dy} F_X(y^{1/3}) \\&= f_X(y^{1/3}) \times \frac{1}{3} y^{-2/3} \quad \text{by chain rule} \\&= \frac{1}{3} y^{-2/3}, \quad 0 \leq y \leq 1 \text{ (and zero otherwise).}\end{aligned}$$

Example 5.1: Solution

$$\begin{aligned}F_Y(y) &= \Pr(Y \leq y) \\&= \Pr(X^3 \leq y) \\&= \Pr(X \leq y^{1/3}) \\&= F_X(y^{1/3}).\end{aligned}$$

Therefore

$$\begin{aligned}f_Y(y) &= \frac{d}{dy} F_X(y^{1/3}) \\&= f_X(y^{1/3}) \times \frac{1}{3} y^{-2/3} \quad \text{by chain rule} \\&= \frac{1}{3} y^{-2/3}, \quad 0 \leq y \leq 1 \text{ (and zero otherwise).}\end{aligned}$$

Example 5.1: Solution

$$\begin{aligned}F_Y(y) &= \Pr(Y \leq y) \\&= \Pr(X^3 \leq y) \\&= \Pr(X \leq y^{1/3}) \\&= F_X(y^{1/3}).\end{aligned}$$

Therefore

$$\begin{aligned}f_Y(y) &= \frac{d}{dy} F_X(y^{1/3}) \\&= f_X(y^{1/3}) \times \frac{1}{3} y^{-2/3} \quad \text{by chain rule} \\&= \frac{1}{3} y^{-2/3}, \quad 0 \leq y \leq 1 \text{ (and zero otherwise).}\end{aligned}$$

Example 5.1: Solution

$$\begin{aligned}F_Y(y) &= \Pr(Y \leq y) \\&= \Pr(X^3 \leq y) \\&= \Pr(X \leq y^{1/3}) \\&= F_X(y^{1/3}).\end{aligned}$$

Therefore

$$\begin{aligned}f_Y(y) &= \frac{d}{dy} F_X(y^{1/3}) \\&= f_X(y^{1/3}) \times \frac{1}{3} y^{-2/3} \quad \text{by chain rule} \\&= \frac{1}{3} y^{-2/3}, \quad 0 \leq y \leq 1 \text{ (and zero otherwise).}\end{aligned}$$

Example 5.1: Solution

$$\begin{aligned}F_Y(y) &= \Pr(Y \leq y) \\&= \Pr(X^3 \leq y) \\&= \Pr(X \leq y^{1/3}) \\&= F_X(y^{1/3}).\end{aligned}$$

Therefore

$$\begin{aligned}f_Y(y) &= \frac{d}{dy} F_X(y^{1/3}) \\&= f_X(y^{1/3}) \times \frac{1}{3} y^{-2/3} \quad \text{by chain rule} \\&= \frac{1}{3} y^{-2/3}, \quad 0 \leq y \leq 1 \text{ (and zero otherwise).}\end{aligned}$$

Example 5.1: Solution

$$\begin{aligned}F_Y(y) &= \Pr(Y \leq y) \\&= \Pr(X^3 \leq y) \\&= \Pr(X \leq y^{1/3}) \\&= F_X(y^{1/3}).\end{aligned}$$


Therefore

$$\begin{aligned}f_Y(y) &= \frac{d}{dy} F_X(y^{1/3}) \\&= f_X(y^{1/3}) \times \frac{1}{3} y^{-2/3} \quad \text{by chain rule} \\&= \frac{1}{3} y^{-2/3}, \quad 0 \leq y \leq 1 \text{ (and zero otherwise).}\end{aligned}$$

Example 5.1: Checking via simulation

To confirm this result, we can simulate Y :

```
1 y = (runif(500))^3
2 hist(y, xlab = 'Y', ylab = 'density', prob=TRUE)
3 curve((1/3)*(x^(-2/3)), from=0, to=1, col="purple", lty
      =2, add=TRUE)
```




Example 5.2

Example

If $X \sim N(0, 1)$, what is the PDF of $Y = X^2$?

Can look at this informally in R:

```
y = (rnorm(1000))^2  
hist(y, xlab="Y", ylab="density", prob=TRUE)
```




Example 5.2

Example

If $X \sim N(0, 1)$, what is the PDF of $Y = X^2$?

Can look at this informally in R:

```
1 y = (rnorm(1000))^2
2 hist(y, xlab="Y", ylab="density", prob=TRUE)
```




Example 5.2

Example

If $X \sim N(0, 1)$, what is the PDF of $Y = X^2$?

Can look at this informally in R:

```
1 y = (rnorm(10000))^2  
2 hist(y, xlab="Y", ylab="density", prob=TRUE)
```




Example 5.2

Example

If $X \sim N(0, 1)$, what is the PDF of $Y = X^2$?

Can look at this informally in R:

```
1 y = (rnorm(1000000))^2  
2 hist(y, xlab="Y", ylab="density", prob=TRUE)
```




Example 5.2

Example

If $X \sim N(0, 1)$, what is the PDF of $Y = X^2$?

Can look at this informally in R:

```
1 y = (rnorm(100000000))^2
2 hist(y, xlab="Y", ylab="density", prob=TRUE)
```



Example 5.2

What about the mathematical solution?

$$\begin{aligned}F_Y(y) &= \Pr(Y \leq y) \\&= \Pr(X^2 \leq y) \\&= \Pr(-\sqrt{y} \leq X \leq \sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \Pr(X \leq -\sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \{1 - \Pr(X \leq \sqrt{y})\} \\&= 2 \cdot \Pr(X \leq \sqrt{y}) - 1 \\&= 2 \cdot F_X(\sqrt{y}) - 1.\end{aligned}$$

Example 5.2

What about the mathematical solution?

$$\begin{aligned}F_Y(y) &= \Pr(Y \leq y) \\&= \Pr(X^2 \leq y) \\&= \Pr(-\sqrt{y} \leq X \leq \sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \Pr(X \leq -\sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \{1 - \Pr(X \leq \sqrt{y})\} \\&= 2 \cdot \Pr(X \leq \sqrt{y}) - 1 \\&= 2 \cdot F_X(\sqrt{y}) - 1.\end{aligned}$$

Example 5.2

What about the mathematical solution?

$$\begin{aligned}F_Y(y) &= \Pr(Y \leq y) \\&= \Pr(X^2 \leq y) \\&= \Pr(-\sqrt{y} \leq X \leq \sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \Pr(X \leq -\sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \{1 - \Pr(X \leq \sqrt{y})\} \\&= 2 \cdot \Pr(X \leq \sqrt{y}) - 1 \\&= 2 \cdot F_X(\sqrt{y}) - 1.\end{aligned}$$

Example 5.2

What about the mathematical solution?

$$\begin{aligned}F_Y(y) &= \Pr(Y \leq y) \\&= \Pr(X^2 \leq y) \\&= \Pr(-\sqrt{y} \leq X \leq \sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \Pr(X \leq -\sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \{1 - \Pr(X \leq \sqrt{y})\} \\&= 2 \cdot \Pr(X \leq \sqrt{y}) - 1 \\&= 2 \cdot F_X(\sqrt{y}) - 1.\end{aligned}$$

Example 5.2

What about the mathematical solution?

$$\begin{aligned}F_Y(y) &= \Pr(Y \leq y) \\&= \Pr(X^2 \leq y) \\&= \Pr(-\sqrt{y} \leq X \leq \sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \Pr(X \leq -\sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \{1 - \Pr(X \leq \sqrt{y})\} \\&= 2 \cdot \Pr(X \leq \sqrt{y}) - 1 \\&= 2 \cdot F_X(\sqrt{y}) - 1.\end{aligned}$$

Example 5.2

What about the mathematical solution?

$$\begin{aligned}F_Y(y) &= \Pr(Y \leq y) \\&= \Pr(X^2 \leq y) \\&= \Pr(-\sqrt{y} \leq X \leq \sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \Pr(X \leq -\sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \{1 - \Pr(X \leq \sqrt{y})\} \\&= 2 \cdot \Pr(X \leq \sqrt{y}) - 1 \\&= 2 \cdot F_X(\sqrt{y}) - 1.\end{aligned}$$

Example 5.2

What about the mathematical solution?

$$\begin{aligned}F_Y(y) &= \Pr(Y \leq y) \\&= \Pr(X^2 \leq y) \\&= \Pr(-\sqrt{y} \leq X \leq \sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \Pr(X \leq -\sqrt{y}) \\&= \Pr(X \leq \sqrt{y}) - \{1 - \Pr(X \leq \sqrt{y})\} \\&= 2 \cdot \Pr(X \leq \sqrt{y}) - 1 \\&= 2 \cdot F_X(\sqrt{y}) - 1.\end{aligned}$$

Example 5.2

Then

$$\begin{aligned}f_Y(y) &= \frac{d}{dy} \left(2 \cdot F_X(y^{1/2}) - 1 \right) \\&= 2f_X(y^{1/2}) \times \frac{1}{2}y^{-1/2} \quad \text{by chain rule} \\&= \frac{1}{\sqrt{y}} f_X(\sqrt{y}) \\&= \frac{1}{\sqrt{y}} \times \frac{1}{\sqrt{2\pi}} \exp \left\{ -\frac{1}{2}(\sqrt{y})^2 \right\} \\&= \frac{1}{\sqrt{2\pi y}} e^{-y/2}, \quad \text{for } y \geq 0, \text{ zero otherwise.}\end{aligned}$$

This is the PDF for a **chi-squared distribution on 1 degree of freedom**.

Example 5.2

Then

$$\begin{aligned}f_Y(y) &= \frac{d}{dy} \left(2 \cdot F_X(y^{1/2}) - 1 \right) \\&= 2f_X(y^{1/2}) \times \frac{1}{2}y^{-1/2} \quad \text{by chain rule} \\&= \frac{1}{\sqrt{y}} f_X(\sqrt{y}) \\&= \frac{1}{\sqrt{y}} \times \frac{1}{\sqrt{2\pi}} \exp \left\{ -\frac{1}{2}(\sqrt{y})^2 \right\} \\&= \frac{1}{\sqrt{2\pi y}} e^{-y/2}, \quad \text{for } y \geq 0, \text{ zero otherwise.}\end{aligned}$$

This is the PDF for a **chi-squared distribution on 1 degree of freedom**.

Example 5.2

Then

$$\begin{aligned}f_Y(y) &= \frac{d}{dy} \left(2 \cdot F_X(y^{1/2}) - 1 \right) \\&= 2f_X(y^{1/2}) \times \frac{1}{2}y^{-1/2} \quad \text{by chain rule} \\&= \frac{1}{\sqrt{y}} f_X(\sqrt{y}) \\&= \frac{1}{\sqrt{y}} \times \frac{1}{\sqrt{2\pi}} \exp \left\{ -\frac{1}{2}(\sqrt{y})^2 \right\} \\&= \frac{1}{\sqrt{2\pi y}} e^{-y/2}, \quad \text{for } y \geq 0, \text{ zero otherwise.}\end{aligned}$$

This is the PDF for a **chi-squared distribution on 1 degree of freedom**.

Example 5.2

Then

$$\begin{aligned}f_Y(y) &= \frac{d}{dy} \left(2 \cdot F_X(y^{1/2}) - 1 \right) \\&= 2f_X(y^{1/2}) \times \frac{1}{2}y^{-1/2} \quad \text{by chain rule} \\&= \frac{1}{\sqrt{y}} f_X(\sqrt{y}) \\&= \frac{1}{\sqrt{y}} \times \frac{1}{\sqrt{2\pi}} \exp \left\{ -\frac{1}{2}(\sqrt{y})^2 \right\} \\&= \frac{1}{\sqrt{2\pi y}} e^{-y/2}, \quad \text{for } y \geq 0, \text{ zero otherwise.}\end{aligned}$$

This is the PDF for a **chi-squared distribution on 1 degree of freedom**.

Example 5.2

Then


$$\begin{aligned}f_Y(y) &= \frac{d}{dy} \left(2 \cdot F_X(y^{1/2}) - 1 \right) \\&= 2f_X(y^{1/2}) \times \frac{1}{2}y^{-1/2} \quad \text{by chain rule} \\&= \frac{1}{\sqrt{y}} f_X(\sqrt{y}) \\&= \frac{1}{\sqrt{y}} \times \frac{1}{\sqrt{2\pi}} \exp \left\{ -\frac{1}{2}(\sqrt{y})^2 \right\} \\&= \frac{1}{\sqrt{2\pi y}} e^{-y/2}, \quad \text{for } y \geq 0, \text{ zero otherwise.}\end{aligned}$$

This is the PDF for a **chi-squared distribution on 1 degree of freedom**.

Example 5.2

To confirm this result we can simulate Y :

```
1 y = (rnorm(1000))^2
2 hist(y, xlab="Y", ylab="density", prob=TRUE)
3 curve((1/sqrt(2*pi*x))*exp(-x/2), from=0, to=10, col="
  purple", lty=2, add=TRUE)
```



Example 5.3

Example

If $X \sim N(\mu, \sigma^2)$, what is the PDF of $Y = \exp(X)$?

This is the example used at the start of this section of the notes.

We say Y has the **log-normal** distribution and write

$Y \sim LN(\mu, \sigma^2)$.

$$F_Y(y) = F_X(\log y) \quad \text{by usual argument with CDF}$$

$$f_Y(y) = \frac{d}{dy} F_X(\log y)$$

$$= f_X(\log y) \times \frac{1}{y} \quad \text{for } y \geq 0$$

$$= \frac{1}{\sigma y \sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{(\log y - \mu)^2}{\sigma^2} \right) \right\}.$$

Example 5.3

Example

If $X \sim N(\mu, \sigma^2)$, what is the PDF of $Y = \exp(X)$?

This is the example used at the start of this section of the notes.

We say Y has the **log-normal** distribution and write $Y \sim LN(\mu, \sigma^2)$.

$$F_Y(y) = F_X(\log y) \quad \text{by usual argument with CDF}$$

$$f_Y(y) = \frac{d}{dy} F_X(\log y)$$

$$= f_X(\log y) \times \frac{1}{y} \quad \text{for } y \geq 0$$

$$= \frac{1}{\sigma y \sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{(\log y - \mu)^2}{\sigma^2} \right) \right\}.$$

Example 5.3

Example

If $X \sim N(\mu, \sigma^2)$, what is the PDF of $Y = \exp(X)$?

This is the example used at the start of this section of the notes.

We say Y has the **log-normal** distribution and write $Y \sim LN(\mu, \sigma^2)$.

$$F_Y(y) = F_X(\log y) \quad \text{by usual argument with CDF}$$

$$f_Y(y) = \frac{d}{dy} F_X(\log y)$$

$$= f_X(\log y) \times \frac{1}{y} \quad \text{for } y \geq 0$$

$$= \frac{1}{\sigma y \sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{(\log y - \mu)^2}{\sigma^2} \right) \right\}.$$

Example 5.3

Example

If $X \sim N(\mu, \sigma^2)$, what is the PDF of $Y = \exp(X)$?

This is the example used at the start of this section of the notes.

We say Y has the **log-normal** distribution and write $Y \sim LN(\mu, \sigma^2)$.

$$F_Y(y) = F_X(\log y) \quad \text{by usual argument with CDF}$$

$$f_Y(y) = \frac{d}{dy} F_X(\log y)$$

$$= f_X(\log y) \times \frac{1}{y} \quad \text{for } y \geq 0$$

$$= \frac{1}{\sigma y \sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{(\log y - \mu)^2}{\sigma^2} \right) \right\}.$$

Example 5.3

Example

If $X \sim N(\mu, \sigma^2)$, what is the PDF of $Y = \exp(X)$?

This is the example used at the start of this section of the notes.

We say Y has the **log-normal** distribution and write $Y \sim LN(\mu, \sigma^2)$.

$$F_Y(y) = F_X(\log y) \quad \text{by usual argument with CDF}$$

$$\begin{aligned} f_Y(y) &= \frac{d}{dy} F_X(\log y) \\ &= f_X(\log y) \times \frac{1}{y} \quad \text{for } y \geq 0 \\ &= \frac{1}{\sigma y \sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{(\log y - \mu)^2}{\sigma^2} \right) \right\}. \end{aligned}$$

5.3 The transformation method

If X has PDF $f_X(x)$ and $h(x)$ is a monotonically increasing or monotonically decreasing function then

- h is invertible and
- the PDF of $Y = h(X)$ is

$$f_Y(y) = f_X(h^{-1}(y)) \times \left| \frac{d(h^{-1}(y))}{dy} \right|$$

when y is in the range of $h(x)$ and zero otherwise.

This result allows direct calculation of $f_Y(y)$ and is called the **univariate transformation method**.

See Lee's scribbles for Proof.

Some Examples

Example

When $X \sim U(0, 1)$, use the transformation method to find the PDF of $Y = X^3$.

$$h(x) = x^3, \quad h^{-1}(y) = y^{1/3}.$$

$$\begin{aligned} f_Y(y) &= 1 \times \frac{d}{dy} y^{1/3} \\ &= \frac{1}{3} y^{-2/3}, \quad 0 \leq y \leq 1. \end{aligned}$$

Some Examples

Example

When $X \sim U(0, 1)$, use the transformation method to find the PDF of $Y = X^3$.

$$h(x) = x^3, \quad h^{-1}(y) = y^{1/3}.$$

$$\begin{aligned} f_Y(y) &= 1 \times \frac{d}{dy} y^{1/3} \\ &= \frac{1}{3} y^{-2/3}, \quad 0 \leq y \leq 1. \end{aligned}$$

Some Examples

Example

When $X \sim U(0, 1)$, use the transformation method to find the PDF of $Y = X^3$.

$$h(x) = x^3, \quad h^{-1}(y) = y^{1/3}.$$

$$\begin{aligned} f_Y(y) &= 1 \times \frac{d}{dy} y^{1/3} \\ &= \frac{1}{3} y^{-2/3}, \quad 0 \leq y \leq 1. \end{aligned}$$

Some Examples

Example

When $X \sim U(0, 1)$, use the transformation method to find the PDF of $Y = X^3$.

$$h(x) = x^3, \quad h^{-1}(y) = y^{1/3}.$$

$$\begin{aligned} f_Y(y) &= 1 \times \frac{d}{dy} y^{1/3} \\ &= \frac{1}{3} y^{-2/3}, \quad 0 \leq y \leq 1. \end{aligned}$$

Some Examples

Example

When $X \sim U(-1, 1)$ and $Y = \exp(X)$, use the transformation method to find the PDF of Y .

$$h(x) = e^x, \quad h^{-1}(y) = \log(y).$$

$$\begin{aligned} f_Y(y) &= \frac{1}{2} \times \left| \frac{d}{dy} \log(y) \right| \\ &= \frac{1}{2|y|} = \frac{1}{2y}, \quad e^{-1} \leq y \leq e^1, \text{ zero otherwise.} \end{aligned}$$

Some Examples

Example

When $X \sim U(-1, 1)$ and $Y = \exp(X)$, use the transformation method to find the PDF of Y .

$$h(x) = e^x, \quad h^{-1}(y) = \log(y).$$

$$\begin{aligned} f_Y(y) &= \frac{1}{2} \times \left| \frac{d}{dy} \log(y) \right| \\ &= \frac{1}{2|y|} = \frac{1}{2y}, \quad e^{-1} \leq y \leq e^1, \text{ zero otherwise.} \end{aligned}$$

Some Examples

Example

When $X \sim U(-1, 1)$ and $Y = \exp(X)$, use the transformation method to find the PDF of Y .

$$h(x) = e^x, \quad h^{-1}(y) = \log(y).$$

$$\begin{aligned} f_Y(y) &= \frac{1}{2} \times \left| \frac{d}{dy} \log(y) \right| \\ &= \frac{1}{2|y|} = \frac{1}{2y}, \quad e^{-1} \leq y \leq e^1, \text{ zero otherwise.} \end{aligned}$$

Some Examples

Example

When $X \sim U(-1, 1)$ and $Y = \exp(X)$, use the transformation method to find the PDF of Y .

$$h(x) = e^x, \quad h^{-1}(y) = \log(y).$$

$$\begin{aligned} f_Y(y) &= \frac{1}{2} \times \left| \frac{d}{dy} \log(y) \right| \\ &= \frac{1}{2|y|} = \frac{1}{2y}, \quad e^{-1} \leq y \leq e^1, \text{ zero otherwise.} \end{aligned}$$

Some Examples

Example

Use the transformation method to find the PDF of the log-normal distribution.

$$h(x) = e^x, \quad h^{-1}(y) = \log(y).$$

$$\begin{aligned} f_Y(y) &= f_X(\log(y)) \times \frac{1}{y} \quad \text{for } y > 0 \\ &= \frac{1}{y} \times \frac{1}{\sigma\sqrt{2\pi}} \exp\left[-\frac{1}{2} \left(\frac{\log(y) - \mu}{\sigma}\right)^2\right]. \end{aligned}$$

Some Examples

Example

Use the transformation method to find the PDF of the log-normal distribution.

$$h(x) = e^x, \quad h^{-1}(y) = \log(y).$$

$$\begin{aligned} f_Y(y) &= f_X(\log(y)) \times \frac{1}{y} \quad \text{for } y > 0 \\ &= \frac{1}{y} \times \frac{1}{\sigma\sqrt{2\pi}} \exp\left[-\frac{1}{2} \left(\frac{\log(y) - \mu}{\sigma}\right)^2\right]. \end{aligned}$$

Some Examples

Example

Use the transformation method to find the PDF of the log-normal distribution.

$$h(x) = e^x, \quad h^{-1}(y) = \log(y).$$

$$f_Y(y) = f_X(\log(y)) \times \frac{1}{y} \quad \text{for } y > 0$$

$$= \frac{1}{y} \times \frac{1}{\sigma\sqrt{2\pi}} \exp\left[-\frac{1}{2} \left(\frac{\log(y) - \mu}{\sigma}\right)^2\right]$$

Some Examples

Example

Use the transformation method to find the PDF of the log-normal distribution.

$$h(x) = e^x, \quad h^{-1}(y) = \log(y).$$

$$\begin{aligned} f_Y(y) &= f_X(\log(y)) \times \frac{1}{y} \quad \text{for } y > 0 \\ &= \frac{1}{y} \times \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{\log(y) - \mu}{\sigma} \right)^2 \right]. \end{aligned}$$

Cauchy example


Example

Suppose $X \sim U(-\frac{\pi}{2}, \frac{\pi}{2})$ and $Y = \tan(X)$.

Use simulation to study the distribution of Y and explicitly compute the PDF.

Solution

```
1 y = tan(runif(100, -pi/2, pi/2))  
2 hist(y, xlab = 'y', ylab = 'frequency', main = '')
```



Solution

This distribution looks odd! Calculate the PDF:

$$h(x) = \tan(x), \quad h^{-1}(y) = \tan^{-1}(y).$$

$$\begin{aligned} f_Y(y) &= \frac{1}{\pi} \frac{d}{dy} (\tan^{-1} y) \quad -\infty < y < \infty, \\ &= \frac{1}{\pi(1+y^2)} \quad -\infty < y < \infty. \end{aligned}$$

This is the **Cauchy distribution**. It has very heavy tails.

Solution

This distribution looks odd! Calculate the PDF:

$$h(x) = \tan(x), \quad h^{-1}(y) = \tan^{-1}(y).$$

$$\begin{aligned} f_Y(y) &= \frac{1}{\pi} \frac{d}{dy} (\tan^{-1} y) \quad -\infty < y < \infty, \\ &= \frac{1}{\pi(1+y^2)} \quad -\infty < y < \infty. \end{aligned}$$

This is the **Cauchy distribution**. It has very heavy tails.

Solution

This distribution looks odd! Calculate the PDF:

$$h(x) = \tan(x), \quad h^{-1}(y) = \tan^{-1}(y).$$

$$\begin{aligned} f_Y(y) &= \frac{1}{\pi} \frac{d}{dy} (\tan^{-1} y) \quad -\infty < y < \infty, \\ &= \frac{1}{\pi(1+y^2)} \quad -\infty < y < \infty. \end{aligned}$$

This is the **Cauchy distribution**. It has very heavy tails.

Solution

This distribution looks odd! Calculate the PDF:

$$h(x) = \tan(x), \quad h^{-1}(y) = \tan^{-1}(y).$$

$$\begin{aligned} f_Y(y) &= \frac{1}{\pi} \frac{d}{dy} (\tan^{-1} y) \quad -\infty < y < \infty, \\ &= \frac{1}{\pi(1+y^2)} \quad -\infty < y < \infty. \end{aligned}$$

This is the Cauchy distribution. It has very heavy tails.

Solution

This distribution looks odd! Calculate the PDF:

$$h(x) = \tan(x), \quad h^{-1}(y) = \tan^{-1}(y).$$

$$\begin{aligned} f_Y(y) &= \frac{1}{\pi} \frac{d}{dy} (\tan^{-1} y) \quad -\infty < y < \infty, \\ &= \frac{1}{\pi(1+y^2)} \quad -\infty < y < \infty. \end{aligned}$$

This is the **Cauchy distribution**. It has *very* heavy tails.

Bonus example

Hourly wind speed maxima (X miles per hour), observed on the roof of Claremont Tower as part of the University's *Urban Observatory*, are assumed exponentially distributed with rate 0.1.

Unfortunately, the data recorded are the logged wind speeds.

Use the transformation method to find the PDF of $Y = \log X$.

Bonus example: Solution

Recall that if $X \sim \text{Exp}(\lambda)$, then it has PDF

$$f_X(x) = \lambda e^{-\lambda x}, \quad x > 0, \quad \lambda > 0.$$

Now here, we have

$$h(x) = \log x, \quad \text{and} \quad h^{-1}(y) = e^y.$$

Therefore

$$\begin{aligned} f_Y(y) &= f_X(h^{-1}(y)) \times \left| \frac{d(h^{-1}(y))}{dy} \right| \\ &= \lambda \exp\{-\lambda \exp\{y\}\} \times \exp\{y\} \\ &= \lambda \exp\{y - \lambda e^y\}, \end{aligned}$$

with $\lambda = 0.1$.

Bonus example: Solution

Recall that if $X \sim \text{Exp}(\lambda)$, then it has PDF

$$f_X(x) = \lambda e^{-\lambda x}, \quad x > 0, \quad \lambda > 0.$$

Now here, we have

$$h(x) = \log x, \quad \text{and} \quad h^{-1}(y) = e^y.$$

Therefore

$$\begin{aligned} f_Y(y) &= f_X(h^{-1}(y)) \times \left| \frac{d(h^{-1}(y))}{dy} \right| \\ &= \lambda \exp\{-\lambda \exp\{y\}\} \times \exp\{y\} \\ &= \lambda \exp\{y - \lambda e^y\}, \end{aligned}$$

with $\lambda = 0.1$.

Bonus example: Solution

Recall that if $X \sim \text{Exp}(\lambda)$, then it has PDF

$$f_X(x) = \lambda e^{-\lambda x}, \quad x > 0, \quad \lambda > 0.$$

Now here, we have

$$h(x) = \log x, \quad \text{and} \quad h^{-1}(y) = e^y.$$

Therefore

$$\begin{aligned} f_Y(y) &= f_X(h^{-1}(y)) \times \left| \frac{d(h^{-1}(y))}{dy} \right| \\ &= \lambda \exp\{-\lambda \exp\{y\}\} \times \exp\{y\} \\ &= \lambda \exp\{y - \lambda e^y\}, \end{aligned}$$

with $\lambda = 0.1$.

Bonus example: Solution

Recall that if $X \sim \text{Exp}(\lambda)$, then it has PDF

$$f_X(x) = \lambda e^{-\lambda x}, \quad x > 0, \quad \lambda > 0.$$

Now here, we have

$$h(x) = \log x, \quad \text{and} \quad h^{-1}(y) = e^y.$$

Therefore

$$\begin{aligned} f_Y(y) &= f_X(h^{-1}(y)) \times \left| \frac{d(h^{-1}(y))}{dy} \right| \\ &= \lambda \exp\{-\lambda \exp\{y\}\} \times \exp\{y\} \\ &= \lambda \exp\{y - \lambda e^y\}, \end{aligned}$$

with $\lambda = 0.1$.

Bonus example: Solution

Recall that if $X \sim \text{Exp}(\lambda)$, then it has PDF

$$f_X(x) = \lambda e^{-\lambda x}, \quad x > 0, \quad \lambda > 0.$$

Now here, we have

$$h(x) = \log x, \quad \text{and} \quad h^{-1}(y) = e^y.$$

Therefore

$$\begin{aligned} f_Y(y) &= f_X(h^{-1}(y)) \times \left| \frac{d(h^{-1}(y))}{dy} \right| \\ &= \lambda \exp\{-\lambda \exp\{y\}\} \times \exp\{y\} \\ &= \lambda \exp\{y - \lambda e^y\}, \end{aligned}$$

with $\lambda = 0.1$.

Bonus example: Solution

Recall that if $X \sim \text{Exp}(\lambda)$, then it has PDF

$$f_X(x) = \lambda e^{-\lambda x}, \quad x > 0, \quad \lambda > 0.$$

Now here, we have

$$h(x) = \log x, \quad \text{and} \quad h^{-1}(y) = e^y.$$

Therefore

$$\begin{aligned} f_Y(y) &= f_X(h^{-1}(y)) \times \left| \frac{d(h^{-1}(y))}{dy} \right| \\ &= \lambda \exp\{-\lambda \exp\{y\}\} \times \exp\{y\} \\ &= \lambda \exp\{y - \lambda e^y\}, \end{aligned}$$

with $\lambda = 0.1$.

Bonus example: Solution

We can check this in R:

```
1 X = rexp(1000, 0.1)
2 Y = log(X)
3 hist(Y, prob=TRUE, main="Y=log(Wind Speeds)", xlab="
  Miles per hour")
4 lambda = 0.1
5 curve(lambda*exp(x-lambda*exp(x)), add=TRUE, col="red",
  lty=2, lwd=2)
```