

INVARIANT DISTRIBUTIONS ON PROJECTIVE SPACES OVER LOCAL FIELDS

GUYAN ROBERTSON

ABSTRACT. Let Γ be an \tilde{A}_n subgroup of $\mathrm{PGL}_{n+1}(\mathbb{K})$, with $n \geq 2$, where \mathbb{K} is a local field with residue field of order q and let $\mathbb{P}_{\mathbb{K}}^n$ be projective n -space over \mathbb{K} . The module of coinvariants $H_0(\Gamma; C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z}))$ is shown to be finite. Consequently there is no nonzero Γ -invariant \mathbb{Z} -valued distribution on $\mathbb{P}_{\mathbb{K}}^n$.

1. INTRODUCTION

Let \mathbb{K} be a nonarchimedean local field with residue field k of order q and uniformizer π . Denote by $\mathbb{P}_{\mathbb{K}}^n$ the set of one dimensional subspaces of the vector space \mathbb{K}^{n+1} , i.e. the set of points in projective n -space over \mathbb{K} . Then $\mathbb{P}_{\mathbb{K}}^n$ is a compact totally disconnected space with the quotient topology inherited from \mathbb{K}^{n+1} , and there is a continuous action of $G = \mathrm{PGL}_{n+1}(\mathbb{K})$ on $\mathbb{P}_{\mathbb{K}}^n$.

Let Γ be a lattice subgroup of G . The abelian group $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})$ of continuous integer-valued functions on $\mathbb{P}_{\mathbb{K}}^n$ has the structure of a Γ -module and the module of coinvariants $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma} = H_0(\Gamma; C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z}))$ is a finitely generated group. Now suppose that Γ is an \tilde{A}_n group [3, 4], i.e. Γ acts freely and transitively on the vertex set of the Bruhat-Tits building of G , which has type \tilde{A}_n . A free group is an \tilde{A}_1 group since it acts freely and transitively on the vertex set of a tree, which is a building of type \tilde{A}_1 . For $n \geq 2$, the \tilde{A}_n groups are unlike free groups. This article proves the following.

Theorem 1.1. *If Γ is an \tilde{A}_n subgroup of $\mathrm{PGL}_{n+1}(\mathbb{K})$, where $n \geq 2$, then $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma}$ is a finite group.*

The proof depends upon the fact that Γ has Kazhdan's property (T). A *distribution* on $\mathbb{P}_{\mathbb{K}}^n$ is a finitely additive \mathbb{Z} -valued measure μ defined on the clopen subsets of $\mathbb{P}_{\mathbb{K}}^n$.

Corollary 1.2. *If Γ is an \tilde{A}_n subgroup of $\mathrm{PGL}_{n+1}(\mathbb{K})$, where $n \geq 2$, then there is no nonzero Γ -invariant \mathbb{Z} -valued distribution on $\mathbb{P}_{\mathbb{K}}^n$.*

This contrasts strongly with the main result of [8] concerning boundary distributions associated with finite graphs. A torsion free lattice subgroup Γ of $\mathrm{PGL}_2(\mathbb{K})$ is a free group, of rank r say. It was shown in [8] that in this case the group of Γ -invariant \mathbb{Z} -valued distributions on $\mathbb{P}_{\mathbb{K}}^1$ is isomorphic to \mathbb{Z}^r . In particular, there are many such distributions.


Date: July 1, 2010.

2000 Mathematics Subject Classification. Primary 20F65, 20G25, 51E24.

Key words and phrases. Buildings, boundary distributions.

2. BACKGROUND

2.1. The Bruhat-Tits building. If \mathbb{K} is a local field, with discrete valuation $v : \mathbb{K}^\times \rightarrow \mathbb{Z}$, let $\mathcal{O} = \{x \in \mathbb{K} : v(x) \geq 0\}$ and let $\pi \in \mathbb{K}$ satisfy $v(\pi) = 1$. A *lattice* L is an \mathcal{O} -submodule of \mathbb{K}^{n+1} of rank $n+1$. In other words $L = \mathcal{O}e_1 + \mathcal{O}e_2 + \cdots + \mathcal{O}e_{n+1}$, for some basis $\{e_1, e_2, \dots, e_{n+1}\}$ of \mathbb{K}^{n+1} . Two lattices L_1 and L_2 are *equivalent* if $L_1 = \alpha L_2$ for some $\alpha \in \mathbb{K}^\times$. The Bruhat-Tits building of $\mathrm{PGL}_{n+1}(\mathbb{K})$ is a two dimensional simplicial complex Δ whose vertices are equivalence classes of lattices in \mathbb{K}^{n+1} [9]. Two lattice classes $[L_0], [L_1]$ are *adjacent* if, for suitable representatives L_1, L_2 , we have $L_0 \subset L_1 \subset \pi^{-1}L_0$. A *simplex* is a set of pairwise adjacent lattice classes. The maximal simplices (*chambers*) are the sets $\{[L_0], [L_1], \dots, [L_n]\}$ where $L_0 \subset L_1 \subset \cdots \subset L_n \subset \pi^{-1}L_0$. These inclusions determine a canonical ordering of the vertices in a chamber, up to cyclic permutation. Each vertex v of Δ has a *type* $\tau(v) \in \mathbb{Z}/(n+1)\mathbb{Z}$, and each chamber of Δ has exactly one vertex of each type. If the Haar measure on \mathbb{K}^{n+1} is normalized so that \mathcal{O}^{n+1} has measure 1 then the type map may be defined by $\tau([L]) = \log_q(\mathrm{vol}(L)) + (n+1)\mathbb{Z}$. The cyclic ordering of the vertices of a chamber coincides with the natural ordering given by the vertex types (Figure 1). Let E^1 denote the set of directed edges $e = (x, y)$ of Δ such that $\tau(y) = \tau(x) + 1$. Write $o(e) = x$ and $t(e) = y$. The subgraph of the 1-skeleton of Δ with edge set E^1 is studied in [5, 7].

FIGURE 1. \tilde{A}_3 case: Cyclic ordering of the vertices of a chamber

Lemma 2.1. *Let C be a chamber of Δ . Then C contains $n+1$ directed edges $e \in E^1$.*

Proof. By [9, Chapter 9.2], there is a basis (e_1, \dots, e_{n+1}) of \mathbb{K}^{n+1} such that the vertices of C are the classes of the lattices

$$\begin{aligned} L_0 &= \pi\mathcal{O}e_1 + \pi\mathcal{O}e_2 + \pi\mathcal{O}e_3 + \cdots + \pi\mathcal{O}e_{n+1} \\ L_1 &= \mathcal{O}e_1 + \pi\mathcal{O}e_2 + \pi\mathcal{O}e_3 + \cdots + \pi\mathcal{O}e_{n+1} \\ L_2 &= \mathcal{O}e_1 + \mathcal{O}e_2 + \pi\mathcal{O}e_3 + \cdots + \pi\mathcal{O}e_{n+1} \\ &\dots\dots\dots \\ L_n &= \mathcal{O}e_1 + \mathcal{O}e_2 + \mathcal{O}e_3 + \cdots + \pi\mathcal{O}e_{n+1}. \end{aligned}$$

Define $L_{n+1} = L_0$. Then the edges C which lie in E^1 are $([L_k], [L_{k+1}])$, where $0 \leq k \leq n$. \square

The building Δ is of type \tilde{A}_n and the action of $\mathrm{GL}_{n+1}(\mathbb{K})$ on the set of lattices induces an action of $\mathrm{PGL}_{n+1}(\mathbb{K})$ on Δ which is transitive on the vertex set. The action of $\mathrm{PGL}_{n+1}(\mathbb{K})$ on Δ is *type rotating* in the sense that, for each $g \in \mathrm{PGL}_{n+1}(\mathbb{K})$, there exists $i \in \mathbb{Z}/(n+1)\mathbb{Z}$ such that $\tau(gv) = \tau(v) + i$ for all vertices $v \in \Delta$.

Fix a vertex $v_0 \in \Delta$ of type 0, and let $\Pi(v_0)$ be the set of vertices adjacent to v_0 . Then $\Pi(v_0)$ has a natural incidence structure: if $u, v \in \Pi(v_0)$ are distinct, then u and v are *incident* if u, v and v_0 lie in a common chamber of Δ . If v_0 is the lattice class $[L_0]$, then $\Pi(v_0)$ consists of the classes $[L]$ where $L_0 \subset L \subset \pi^{-1}L_0$, and one can associate to $[L] \in \Pi(v_0)$ the subspace $v = L/L_0$ of $\pi^{-1}L_0/L_0 \cong k^{n+1}$. Thus we may identify $\Pi(v_0)$ with the flag complex of subspaces of the vector space k^{n+1} . Under this identification, a vertex $v \in \Pi(v_0)$ has type $\tau(v) = \dim(v) + \mathbb{Z}/(n+1)\mathbb{Z}$ where $\dim(v)$ is the dimension of v over k . A chamber C of Δ which contains v_0 has vertices v_0, v_1, \dots, v_n where $(0) = v_0 \subset v_1 \subset \dots \subset v_n \subset k^{n+1}$ is a complete flag. For brevity, write $C = \{v_0 \subset v_1 \subset \dots \subset v_n\}$.

Definition 2.2. If $e = ([L_0], [L_1]) \in E^1$, where $L_0 \subset L_1 \subset \pi^{-1}L_0$ and $\tau([L_1]) = \tau([L_0]) + 1$, then define $\Omega(e)$ to be the set of lines $\ell \in \mathbb{P}_{\mathbb{K}}^n$ such that $L_1 = L_0 + (\ell \cap \pi^{-1}L_0)$. The sets $\Omega(e)$, $e \in E^1$, form a basis for the topology on $\mathbb{P}_{\mathbb{K}}^n$ (c.f. [10, Ch.II.1.1], [1, 1.6]).

Lemma 2.3. If $e \in E^1$, then $\Omega(e)$ may be expressed as a disjoint union of q^n sets

$$(1) \quad \Omega(e) = \bigsqcup_{\substack{o(e')=t(e) \\ \Omega(e') \subset \Omega(e)}} \Omega(e').$$

Proof. Let $e = ([L_0], [L_1]) \in E^1$, where $L_0 \subset L_1 \subset \pi^{-1}L_0$ and $\tau([L_1]) = \tau([L_0]) + 1$. If $\ell \in \Omega(e)$ then $L_1 = L_0 + (\ell \cap \pi^{-1}L_0)$. Choose $e' = ([L_1], [L_2])$ where $L_2 = L_0 + (\ell \cap \pi^{-2}L_0)$. Now $L_0 \subset L_1 \subset L_2 \subset \pi^{-1}L_1$ and L_2/L_1 is a 1-dimensional subspace of $\pi^{-1}L_1/L_1 \cong k^{n+1}$. Moreover, L_2/L_1 is not incident with the n -dimensional subspace $\pi^{-1}L_0/L_1$ of $\pi^{-1}L_1/L_1 \cong k^{n+1}$. There are precisely q^n such 1-dimensional subspaces of k^{n+1} , each of which corresponds to an edge $e' \in E^1$. \square

Lemma 2.4. If ξ is a fixed vertex of Δ , then $\mathbb{P}_{\mathbb{K}}^n$ may be expressed as a disjoint union

$$(2) \quad \mathbb{P}_{\mathbb{K}}^n = \bigsqcup_{o(e)=\xi} \Omega(e).$$

Proof. Let $\xi = [L_0]$, where L_0 is a lattice. If $\ell \in \mathbb{P}_{\mathbb{K}}^n$, define the lattice $L_1 = L_0 + (\ell \cap \pi^{-1}L_0)$. Then $L_0 \subset L_1 \subset \pi^{-1}L_0$ and $\tau([L_1]) = \tau([L_0]) + 1$, since L_0 is maximal in L_1 . Thus the edge $e = ([L_0], [L_1])$ lies in E^1 , and $\ell \in \Omega(e)$. \square

Lemma 2.5. Let C be a chamber of Δ and denote the directed edges of $C \cap E^1$ by e_0, e_1, \dots, e_n . Then $\mathbb{P}_{\mathbb{K}}^n$ may be expressed as a disjoint union

$$(3) \quad \mathbb{P}_{\mathbb{K}}^n = \bigsqcup_{i=0}^n \Omega(e_i).$$

Proof. Let C have vertex set $\{[L_0], [L_1], \dots, [L_n]\}$ where $L_0 \subset L_1 \subset \dots \subset L_n \subset \pi^{-1}L_0$. Let $\ell = \mathbb{K}a \in \mathbb{P}_{\mathbb{K}}^n$, where $a \in \mathbb{K}^{n+1}$ is scaled so that $a \in \pi^{-1}L_0 - L_0$. Then $a \in L_{i+1} - L_i$ for some i , where $L_{i+1}/L_i \cong k$ and $L_{n+1} = \pi^{-1}L_0$. Thus $\ell \in \Omega(e_i)$. \square

2.2. \tilde{A}_n groups. From now on let $\Pi = \Pi(v_0)$, the set of neighbours of the fixed vertex $v_0 \in \Delta$. Thus Π is isomorphic to the flag complex of subspaces of k^{n+1} and a chamber C of Δ which contains v_0 is a complete flag $\{v_0 \subset v_1 \subset \cdots \subset v_n\}$. For $1 \leq r \leq n$, let $\Pi_r = \{u \in \Pi(v_0) : \dim u = r\}$.

Now suppose that Γ is an \tilde{A}_n group i.e. Γ acts freely and transitively on the vertex set of Δ [3, 4]. Then for each $v \in \Pi(v_0)$, there is a unique element $g_v \in \Gamma$ such that $g_v v_0 = v$. If $v \in \Pi(v_0)$, then $g_v^{-1} v_0$ also lies in $\Pi(v_0)$, and $\lambda(v) = g_v^{-1} v_0$ defines an involution $\lambda : \Pi(v_0) \rightarrow \Pi(v_0)$ such that $g_{\lambda(v)} = g_v^{-1}$. Let $\mathcal{T} = \{(u, v, w) \in \Pi(v_0)^3 : g_u g_v g_w = 1\}$. If $(u, v, w) \in \mathcal{T}$ then w is uniquely determined by (u, v) and there is a bijective correspondence between triples $(u, v, w) \in \mathcal{T}$ and directed triangles $(v_0, \lambda(u), v)$ of Δ containing v_0 . By [6, Proposition 2.2], the abstract group Γ has a presentation with generating set $\{g_v : v \in \Pi(v_0)\}$ and relations

$$(4a) \quad g_u g_{\lambda(u)} = 1, \quad u \in \Pi(v_0);$$

$$(4b) \quad g_u g_v g_w = 1, \quad (u, v, w) \in \mathcal{T}.$$

If $u \in \Pi(v_0)$ then $\tau(g_u v_0) = \tau(u) = \tau(u) + \tau(v_0)$. Hence $\tau(g_u x) = \tau(u) + \tau(x)$ for each vertex x of Δ , since g_u is type rotating. In particular, if $u, v \in \Pi(v_0)$ then

$$(5) \quad \tau(g_u g_v v_0) = \tau(u) + \tau(v).$$

It follows from (5) that

$$\tau(\lambda(u)) = -\tau(u)$$

for each $u \in \Pi$. Also, if $(u, v, w) \in \mathcal{T}$, then

$$\tau(u) + \tau(v) + \tau(w) = 0.$$

Let $C = \{v_0 \subset v_1 \subset \cdots \subset v_n\}$ be a chamber of Δ containing v_0 . Since the vertices v_{i-1} and v_i are adjacent, so are the vertices $v_0 = g_{v_{i-1}}^{-1} v_{i-1}$ and $g_{v_{i-1}}^{-1} g_{v_i} v_0 = g_{v_{i-1}}^{-1} v_i$. Also $\tau(g_{v_{i-1}}^{-1} g_{v_i} v_0) = \tau(v_i) - \tau(v_{i-1}) = 1$. Therefore $g_{v_{i-1}}^{-1} g_{v_i} = g_{a_i}$ where $a_i \in \Pi_1$, $v_{n+1} = v_0$ and $g_{v_0} = 1$. Thus $g_{a_1} g_{a_2} \cdots g_{a_k} = g_{v_k}$ ($1 \leq k \leq n$) and $g_{a_1} g_{a_2} \cdots g_{a_{n+1}} = 1$.

The $(n+1)$ -tuple $\sigma(C) = (a_1, a_2, \dots, a_{n+1}) \in \Pi_1^{n+1}$ is uniquely determined by the chamber C containing v_0 . Denote by \mathfrak{S} the set of all $(n+1)$ -tuples $\sigma(C)$ associated with such chambers C . If $u \in \Pi(v_0)$ with $\dim(u) = k$, then u is a vertex of a chamber C containing v_0 . Therefore

$$(6) \quad g_u = g_{a_1} g_{a_2} \cdots g_{a_k}, \quad \text{where } a_i \in \Pi_1, 1 \leq i \leq k.$$

In particular, the set $\{g_a : a \in \Pi_1\}$ generates Γ . Since $g_{\lambda(u)} = g_u^{-1}$, we have

$$(7) \quad g_{\lambda(u)} = g_{a_{i+1}} \cdots g_{a_{n+1}}.$$

Note that the expression (6) for g_u is not unique, but depends on the choice of the chamber C containing u and v_0 . An edge in E^1 has the form $(x, g_a x)$ where $a \in \Pi_1$.

Lemma 2.6. *The \tilde{A}_n group Γ has a presentation with generating set $\{g_a : a \in \Pi_1\}$ and relations*

$$(8) \quad g_{a_1} g_{a_2} \cdots g_{a_{n+1}} = 1, \quad (a_1, a_2, \dots, a_{n+1}) \in \mathfrak{S}.$$

Proof. It is enough to show that the relations (4) follow from the relations (8). Let $(u, v, w) \in \mathcal{T}$ with $\dim(u) = i, \dim v = j$ and $\dim w = k$, where $i + j + k \equiv 0 \pmod{n+1}$. Choose a chamber $C = \{v_0 \subset v_1 \subset \cdots \subset v_n\}$ containing $\{v_0, g_u v_0, g_u g_v v_0\}$. Let $(a_1, a_2, \dots, a_{n+1}) = \sigma(C) \in \Pi_1^{n+1}$ be the element of \mathfrak{S} determined by C . Then $g_u v_0$ is the vertex of C of type i , so $g_u = g_{a_1} g_{a_2} \cdots g_{a_i}$.

Suppose that $j < n + 1 - i$. Then $g_u g_v v_0$ is the vertex of C of type $i + j$ and $g_u g_v = g_{a_1} g_{a_2} \cdots g_{a_{i+j}}$. Thus $g_v = g_{a_{i+1}} \cdots g_{a_{i+j}}$ and $g_w = g_{a_{i+j+1}} \cdots g_{a_{n+1}}$. Therefore

$$g_u g_v g_w = g_{a_1} g_{a_2} \cdots g_{a_{n+1}}.$$

Suppose that $j > n + 1 - i$. Then $g_u g_v v_0$ has type $i + j - n - 1$ and

$$g_u g_v = g_{a_1} g_{a_2} \cdots g_{a_{i+j-n-1}} = g_{a_1} g_{a_2} \cdots g_{a_{n+1}} g_{a_1} \cdots g_{a_{i+j-n-1}}.$$

Thus $g_v = g_{a_{i+1}} \cdots g_{a_{n+1}} g_{a_1} \cdots g_{a_{i+j-n-1}}$ and $g_w = g_{a_{i+j-n}} \cdots g_{a_{n+1}}$. Therefore

$$g_u g_v g_w = (g_{a_1} g_{a_2} \cdots g_{a_{n+1}})^2.$$

In each case the relations (4b) follow from the relations (8). The same is true for the relations (4a), by equation (7). \square

3. THE COINVARIANTS

If Γ is an \tilde{A}_n group acting on Δ , then Γ acts on $\mathbb{P}_{\mathbb{K}}^n$, and the abelian group $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})$ has the structure of a Γ -module, with $(g \cdot f)(\ell) = f(g^{-1}\ell)$, $g \in \Gamma$, $\ell \in \mathbb{P}_{\mathbb{K}}^n$. The module of coinvariants, $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma}$, is the quotient of $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})$ by the submodule generated by $\{g \cdot f - f : g \in \Gamma, f \in C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})\}$. If $f \in C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})$ then let $[f]$ denote its class in $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma}$. Also, let $\mathbf{1}$ denote the constant function defined by $\mathbf{1}(\ell) = 1$ for $\ell \in \mathbb{P}_{\mathbb{K}}^n$, and let $\varepsilon = [\mathbf{1}]$.

If $e \in E^1$, let χ_e be the characteristic function of $\Omega(e)$. For each $g \in \Gamma$, the functions χ_e and $g \cdot \chi_e = \chi_{ge}$ project to the same element in $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma}$. Any edge $e \in E^1$ is in the Γ -orbit of some edge $(v_0, g_a v_0)$, where $a \in \Pi_1$ is uniquely determined by e . Therefore it makes sense to denote by $[a]$ the class of χ_e in $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma}$.

Lemma 3.1. *The group $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma}$ is finitely generated, with generating set $\{[a] : a \in \Pi_1\}$.*

Proof. Every clopen set V in $\mathbb{P}_{\mathbb{K}}^n$ may be expressed as a finite disjoint union of sets of the form $\Omega(e)$, $e \in E^1$. Any function $f \in C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})$ is bounded, by compactness of $\mathbb{P}_{\mathbb{K}}^n$, and so takes finitely many values $n_i \in \mathbb{Z}$. Therefore f may be expressed as a finite sum $f = \sum_j n_j \chi_{e_j}$, with $e_j \in E^1$. The result follows, since $\{\chi_e : e \in E^1\} = \{[a] : a \in \Pi_1\}$. \square

Suppose that $e, e' \in E^1$ with $o(e') = t(e) = x$, so that $o(e) = g_{\lambda(a)}x$ and $t(e') = g_b x$ for (unique) $a, b \in \Pi_1$. Then, by the proof of Lemma 2.3, $\Omega(e') \subset \Omega(e)$ if and only if $b \cap \lambda(a) = (0)$.

Equations (1) and (2) imply the following relations in $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma}$.

$$(9a) \quad \varepsilon = \sum_{a \in \Pi_1} [a];$$

$$(9b) \quad [a] = \sum_{\substack{b \in \Pi_1 \\ b \cap \lambda(a) = (0)}} [b], \quad a \in \Pi_1.$$

It is easy to see that $|\Pi_1| = \frac{q^{n+1}-1}{q-1}$. If $a \in \Pi_1$, then $\lambda(a) \in \Pi_n$ and so the number of elements $b \in \Pi_1$ which are incident with $\lambda(a)$ is $\frac{q^n-1}{q-1}$. Thus there exist q^n elements $b \in \Pi_1$ such that $b \cap \lambda(a) = (0)$. In other words, the right side of (9b) contains q^n terms. As a first step towards proving that $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma}$ is finite, we show that the element $\varepsilon = [\mathbf{1}]$ has finite order.

Lemma 3.2. *In the group $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma}$, $(q^n - 1)\varepsilon = 0$.*

Proof. By (9a) and (9b),

$$\varepsilon = \sum_{a \in \Pi_1} [a] = \sum_{a \in \Pi_1} \left(\sum_{\substack{b \in \Pi_1 \\ b \cap \lambda(a) = (0)}} [b] \right) = \sum_{b \in \Pi_1} q^n [b] = q^n \varepsilon.$$

□

We can now prove Theorem 1.1. It follows from (3) that if $(a_1, a_2, \dots, a_{n+1}) \in \mathfrak{S}$ then

$$(10) \quad \sum_{i=1}^{n+1} [a_i] = \varepsilon.$$

Therefore, by Lemmas 2.6 and 3.1, there is a homomorphism θ from Γ onto the abelian group $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma} / \langle \varepsilon \rangle$ defined by $\theta(g_a) = [a] + \langle \varepsilon \rangle$, for $a \in \Pi_1$.

The \tilde{A}_n group Γ has Kazhdan's property (T) [2, Theorems 1.6.1 and 1.7.1]. It follows that $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma} / \langle \varepsilon \rangle$ is finite [2, Corollary 1.3.5]. Therefore $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma}$ is also finite, since $\langle \varepsilon \rangle$ is finite, by Lemma 3.2. □

Distributions. A *distribution* on $\mathbb{P}_{\mathbb{K}}^n$ is a finitely additive \mathbb{Z} -valued measure μ defined on the clopen subsets of $\mathbb{P}_{\mathbb{K}}^n$ [1, 1.4]. By integration, a distribution may be regarded as a \mathbb{Z} -linear function on the group $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})$. Therefore a Γ -invariant distribution defines a homomorphism $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma} \rightarrow \mathbb{Z}$. This homomorphism is necessarily trivial, since $C(\mathbb{P}_{\mathbb{K}}^n, \mathbb{Z})_{\Gamma}$ is finite. This proves Corollary 1.2.

REFERENCES

- [1] G. Alon and E. de Shalit, On the cohomology of Drinfel'd's p -adic symmetric domain, *Israel J. Math.* **129** (2002), 1–20.
- [2] B. Bekka, P. de la Harpe and A. Valette, *Kazhdan's Property (T)*, Cambridge University Press, Cambridge, 2008.
- [3] D. I. Cartwright, Groups acting simply transitively on the vertices of a building of type \tilde{A}_n , *Groups of Lie type and their Geometries*, W. M. Kantor and L. Di Martino, editors, 43–76. Cambridge University Press, 1995.
- [4] D. I. Cartwright and T. Steger, A family of \tilde{A}_n groups, *Israel J. Math.* **103** (1998), 125–140.
- [5] D. I. Cartwright, P. Solé and A. Žuk, Ramanujan geometries of type \tilde{A}_n , *Discrete Math.* **269** (2003), 35–43.
- [6] D. I. Cartwright, A. M. Mantero, T. Steger and A. Zappa, Groups acting simply transitively on the vertices of a building of type \tilde{A}_2 , I, *Geom. Ded.* **47** (1993), 143–166.
- [7] A. Lubotzky, B. Samuels and U. Vishne, Ramanujan complexes of type \tilde{A}_d , *Israel J. Math.* **149** (2005), 267–299.
- [8] G. Robertson, Invariant boundary distributions associated with finite graphs, *J. Combin. Theory Ser. A*, **115** (2008), 1272–1278.
- [9] M. Ronan, *Lectures on Buildings*, University of Chicago Press, 2009.
- [10] J.-P. Serre, *Trees*, Springer-Verlag, Berlin, 1980.

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEWCASTLE, NE1 7RU, ENGLAND, U.K.

E-mail address: a.g.robertson@ncl.ac.uk