

MAS8304

NEWCASTLE UNIVERSITY

SCHOOL OF MATHEMATICS & STATISTICS

---

SEMESTER 2 2012/2013

---

MAS8304

Environmental Extremes: Mid-semester test

Time allowed: 50 minutes

*Candidates should attempt all questions. Marks for each question are indicated.*

*There are THREE questions on this paper.*

*Answers to questions should be entered directly on this question paper in the spaces provided.  
This question paper must be handed in at the end of the test.*

**Name:**.....

**Exponential distribution:**

If  $X \sim \text{Exp}(\lambda)$ , then it has distribution function

$$F_X(x; \lambda) = 1 - e^{-\lambda x},$$

$$x > 0, \lambda > 0.$$

**Generalised Extreme Value distribution:**

If  $X \sim \text{GEV}(\mu, \sigma, \xi)$  then it has distribution function

$$G_X(x; \mu, \sigma, \xi) = \begin{cases} \exp \left\{ - \left[ 1 + \xi \left( \frac{x - \mu}{\sigma} \right) \right]_+^{-1/\xi} \right\}, & \xi \neq 0; \\ \exp \left\{ - \exp \left[ - \left( \frac{x - \mu}{\sigma} \right) \right] \right\}, & \xi = 0, \end{cases}$$

$$-\infty < \mu < \infty, -\infty < \xi < \infty, \sigma > 0, a_+ = \max(0, a).$$

**Generalised Pareto distribution:**

If  $X \sim \text{GPD}(\tilde{\sigma}, \xi)$  then it has distribution function

$$H_X(x; \tilde{\sigma}, \xi) = \begin{cases} 1 - \left( 1 + \frac{\xi x}{\tilde{\sigma}} \right)_+^{-1/\xi}, & \xi \neq 0; \\ 1 - \exp \left( -\frac{x}{\tilde{\sigma}} \right) & \xi = 0, \end{cases}$$

$$-\infty < \xi < \infty, \tilde{\sigma} > 0, a_+ = \max(0, a).$$

1. Suppose  $X_1, \dots, X_n$  is a sequence of independent Weibull random variables, that is

$$F(x) = \exp \{ -(-x)^\alpha \}.$$

Suppose  $\alpha = 1$ . Show that the limit distribution of  $(M_n - b_n)/a_n$ , where

$$M_n = \max(X_1, \dots, X_n),$$

is of extreme value type, and identify the distribution. Also clearly identify the values of the normalising constants  $a_n$  and  $b_n$ .

**Answer:**

[Total Q1: 10 marks]

2. Annual maximum flood discharges, in units of 1000 cubed feet per second, are available for two locations on the Fox River in Wisconsin, USA: Berlin (upstream) and Wrightstown (downstream). The data span a period of 33 years, from 1980 to 2012 (inclusive). Two analysts work separately on the data collected at Berlin and Wrightstown.

Lucy, the analyst working on the flood data at Berlin, assumes a GEV with  $\xi = 0$ , as is often the case for extreme river discharges at upstream locations. She finds the following maximum likelihood estimates of the remaining GEV location and scale parameters:

$$\hat{\mu} = 3.38 \quad \text{and} \quad \hat{\sigma} = 1.45,$$

with associated hessian of the corresponding log-likelihood function:

$$H = \begin{pmatrix} -11.771 & 4.176 \\ 4.176 & -19.864 \end{pmatrix}.$$

Duncan, the analyst working on the flood data for Wrightstown, assumes a GEV with  $\xi \neq 0$ , and uses the `ismev` package in R. After storing the data in `foxflood`, he obtains the following output:

```
A=gev.fit(foxflood)
$conv
[1] 0

$nllh
[1] 98.01564

$mle
[1] 6.0169817 5.1352751 -0.4485899

$se
[1] 1.0287297 0.8266614 0.1744666
```

- (a) Why does the R output above support Duncan's decision to assume that  $\xi \neq 0$  for Wrightstown? Comment on the tail behaviour of the flood discharges at Wrightstown, as suggested by Alan's results.

[4 marks]

**Answer:**

- (b) Use Lucy's results to obtain estimated standard errors for  $\hat{\mu}$  and  $\hat{\sigma}$  at Berlin.  
[6 marks]

**Answer:**

- (c) By constructing confidence intervals for the GEV location parameters in the usual way, comment on whether or not there is a significant difference between the annual maximum flood discharges at these two locations on the Fox river.  
[6 marks]

**Answer:**

- (d) For both locations, find the probability that, this year, the annual maximum flood discharge will exceed  $5000 \text{ feet}^3/\text{s}$ .

[6 marks]

**Answer:**

- (e) Over the next 100 years, in how many years can we expect the Fox river to have a flood discharge in excess of  $5000 \text{ feet}^3/\text{s}$  at *both* locations? [*Hint: Assume the river floods independently at Berlin and Wrightstown*]

[3 marks]

**Answer:**

[Total Q1: 25 marks]

3. You are working with a team of climatologists who are interested in temperature patterns across continental Europe. Daily minimum temperatures ( $^{\circ}\text{C}$ ) are recorded at Sarajevo, in Bosnia–Herzegovina, between 2006–2011 (inclusive). To work with the standard extreme value models for extremely *large* observations (as opposed to the extremely *small* observations as you have here), you decide to negate your data.

- (a) In the context of the generalised extreme value distribution, what effect does working with the negated data have on estimates of the GEV parameters?

[2 marks]

**Answer:**

- (b) You decide to perform an analysis of threshold exceedances. With reference to Figure 1 overleaf, briefly explain why you might identify extremes as values which exceed a threshold of  $u = -10^{\circ}\text{C}$ .

[1 mark]

**Answer:**

- (c) You analyse the data in R. The negated temperatures are stored in the vector `neg.sarajevo`; you then use the `ismev` function `gpd.fit`, resulting in the (edited) output shown below. Look at it, and then answer the following questions.

```
B=gpd.fit(neg.sarajevo,-10)
$threshold
[1] -10
```

```
$nexc
[1] 233
```

```
$conv
[1] 0
```

```
$nllh
[1] 636.7434
```

```
$mle
[1] 7.4306027 -0.2728779
```

```
$se
[1] 0.59746049 0.04995889
```

- (i) Estimate the threshold exceedance rate  $\lambda$ .

[2 marks]

**Answer:**

- (ii) Given that  $\text{cov}(\hat{\sigma}, \hat{\xi}) = -0.025$ , complete the variance–covariance matrix  $V$  for the parameter vector  $\boldsymbol{\theta} = (\lambda, \sigma, \xi)^T$ .

[5 marks]

**Answer:**

$$V = \begin{pmatrix} & & \\ & & \\ & & \end{pmatrix}$$

- (d) With reference to Figure 1 overleaf, find the estimated 50–year return level *minimum* daily temperature for Sarajevo, with it's associated 95% profile log–likelihood confidence interval. [Hint:  $\chi_1^2(0.05) = 3.841$ ]

[5 marks]

**Answer:**

[Total Q3: 15 marks]

**THE END**






Figure 1: Mean residual life plot (top); profile log-likelihood curve for the 50-year return level.