

Chapter 9


Continuous Probability Models

Outline

- Probability Density Functions (pdfs)
- The Uniform Distribution
- The Exponential Distribution

Continuous Data

Histogram of 10 obsns, 100 obsns, 1,000 obsns, 10,000 obsns



Probability Density Functions (pdfs)

The key features of pdfs are

1. pdfs never take negative values
2. the area under a pdf is one: $P(-\infty < X < \infty) = 1$
3. areas under the curve correspond to probabilities
4. $P(X \leq x) = P(X < x)$ since $P(X = x) = 0$.


The Uniform Distribution

- Outcomes measured on a continuous scale.
- All outcomes equally likely.

Example: Environmental Health Officers visit local hotel at a “random” time during the working day (9.00 to 18.00)

- Let X be the time to their arrival at the hotel measured in terms of minutes from the start of the day.
- Then X is a random variable with a uniform distribution between 0 and 540, with pdf

$$f(x) = \begin{cases} \frac{1}{540} & \text{for } 0 \leq x \leq 540 \\ 0 & \text{otherwise.} \end{cases}$$



The Uniform Distribution

General case:

Random variable X with a uniform distribution on a to b has pdf

$$f(x) = \begin{cases} \frac{1}{b-a} & \text{for } a \leq x \leq b \\ 0 & \text{otherwise} \end{cases}$$

and probabilities can be calculated using the formula

$$P(X \leq x) = \begin{cases} 0 & \text{for } x < a \\ \frac{x-a}{b-a} & \text{for } a \leq x \leq b \\ 1 & \text{for } x > b. \end{cases}$$

Probability that the inspectors visit the hotel in the morning (within 180 minutes after 9am) is

$$P(X \leq 180) = \frac{180 - 0}{540 - 0} = \frac{1}{3}$$

Probability of a visit during the lunch hour (12.30 to 13.30) is

$$\begin{aligned} P(210 \leq X \leq 270) &= P(X \leq 270) - P(X < 210) \\ &= \frac{270 - 0}{540 - 0} - \frac{210 - 0}{540 - 0} \\ &= \frac{270 - 210}{540} \\ &= \frac{60}{540} \\ &= \frac{1}{9}. \end{aligned}$$

The Uniform Distribution

Mean:

$$E(X) = \mu = \frac{(a + b)}{2}$$

Variance:

$$Var(X) = \sigma^2 = \frac{(b - a)^2}{12}$$

The Exponential Distribution

- used to model lifetimes of products and times between “random” events
- arrivals of orders, customers in a queueing system, ...
- has one (positive) parameter λ


The Exponential Distribution

General form of pdf

$$f(x) = \begin{cases} \lambda e^{-\lambda x} & \text{for } x \geq 0, \\ 0 & \text{otherwise} \end{cases}$$

and probabilities can be calculated using

$$P(X \leq x) = \begin{cases} 0 & \text{for } x < 0 \\ 1 - e^{-\lambda x} & \text{for } x > 0. \end{cases}$$



The Exponential Distribution

Main features

1. it refers to positive quantities: $x > 0$
2. larger values of x are increasingly unlikely – exponential decay
3. the value of λ fixes the rate of decay – larger values of λ correspond to more rapid decay.

The Exponential Distribution

- X = time between use of pay phone
- Model using exponential distribution with $\lambda = 0.3$

Probability of the gap between phone users being less than 5 minutes is

$$P(X < 5) = 1 - e^{-0.3 \times 5} = 1 - 0.223 = 0.777.$$

Probability that the gap is more than 10 minutes is

$$\begin{aligned}P(X > 10) &= 1 - P(X \leq 10) \\&= 1 - (1 - e^{-0.3 \times 10}) \\&= e^{-0.3 \times 10} \\&= 0.050\end{aligned}$$

Probability that the gap is between 5 and 10 minutes is

$$\begin{aligned}P(5 < X < 10) &= P(X < 10) - P(X \leq 5) \\&= 0.950 - 0.777 \\&= 0.173\end{aligned}$$

The Exponential Distribution

Mean:

$$E(X) = \mu = \frac{1}{\lambda}$$

Variance:

$$Var(X) = \sigma^2 = \frac{1}{\lambda^2}$$