




Characterising Jupiter's dynamo radius using its magnetic energy spectrum

Yue-Kin Tsang

School of Mathematics, University of Leeds

Chris Jones (*Leeds*)

Let's start on Earth...



- **core-mantle boundary (CMB):** *sharp boundary* between the **non-conducting mantle** and the **conducting outer core**
⇒ dynamo action entirely confined within the outer core
- **dynamo radius r_{dyn} :** top of the dynamo region $\approx r_{\text{cmb}}$
- one way to deduce r_{cmb} from *observation at the surface*: **magnetic energy spectrum**

Gauss coefficients g_{lm} and h_{lm}


- Outside the dynamo region, $r > r_{\text{dyn}}$:

$$j = 0$$

$$\nabla \times \mathbf{B} = \mu_0 \mathbf{j} = \mathbf{0} \implies \mathbf{B} = -\nabla \Psi$$

$$\nabla \cdot \mathbf{B} = 0 \implies \nabla^2 \Psi = 0$$

$a = \text{radius of Earth}$



- Consider only internal sources,

$$\Psi(r, \theta, \phi) = a \sum_{l=1}^{\infty} \sum_{m=0}^l \left(\frac{a}{r}\right)^{l+1} \hat{P}_{lm}(\cos \theta) (g_{lm} \cos m\phi + h_{lm} \sin m\phi)$$

\hat{P}_{lm} : Schmidt's semi-normalised associated Legendre polynomials

- g_{lm} and h_{lm} can be determined from magnetic field measured at the planetary surface ($r \approx a$)

The Lowes spectrum

- Average magnetic energy over a spherical surface of radius r

$$E_B(r) = \frac{1}{2\mu_0} \frac{1}{4\pi} \oint |\mathbf{B}(r, \theta, \phi)|^2 \sin \theta \, d\theta \, d\phi$$



- Inside the source-free region $r_{\text{dyn}} < r < a$,

$$2\mu_0 E_B(r) = \sum_{l=1}^{\infty} \left[\left(\frac{a}{r} \right)^{2l+4} (l+1) \sum_{m=0}^l (g_{lm}^2 + h_{lm}^2) \right]$$

- **Lowes spectrum** (magnetic energy as a function of l):

$$\begin{aligned} R_l(r) &= \left(\frac{a}{r} \right)^{2l+4} (l+1) \sum_{m=0}^l (g_{lm}^2 + h_{lm}^2) \\ &= \left(\frac{a}{r} \right)^{2l+4} R_l(a) \quad (\text{downward continuation}) \end{aligned}$$

Estimate location of CMB using the Lowes spectrum



- downward continuation from a to r_{cmb} through the mantle ($j = 0$):


$$\ln R_l(a) = 2 \ln \left(\frac{r_{\text{cmb}}}{a} \right) l + 4 \ln \left(\frac{r_{\text{cmb}}}{a} \right) + \ln R_l(r_{\text{cmb}})$$

- white source hypothesis:** turbulence in the core leads to an *even distribution of magnetic energy* across different scales l ,


$R_l(r_{\text{cmb}})$ is independent of l

- $r_{\text{cmb}} \approx 0.55a \approx 3486 \text{ km}$ agrees very well with results from seismic waves observations

Interior structure of Jupiter



(NASA JPL)




theoretical $\sigma(r)$ (French et al. 2012)

- low temperature and pressure near surface \Rightarrow gaseous molecular H/He
- extremely high temperature and pressure inside \Rightarrow liquid metallic H
- core?
- transition from molecular to metallic hydrogen is continuous
- conductivity $\sigma(r)$ varies smoothly with radius r

At what depth does dynamo action start?


Lowes spectrum from the Juno mission



(Connerney et al. 2018)

- Juno's spacecraft reached Jupiter on 4th July, 2016
- currently in a 53-day orbit, measuring Jupiter's magnetic field (and other data)
- $R_l(r_J)$ up to $l = 10$ from latest measurement (8 flybys)
- Lowes' radius: $r_{\text{lowes}} \approx 0.85 r_J$ ($r_J = 6.9894 \times 10^7 \text{m}$)

Lowes spectrum from the Juno mission



(Connerney et al. 2018)

- Juno's spacecraft reached Jupiter on 4th July, 2016
- currently in a 53-day orbit, measuring Jupiter's magnetic field (and other data)
- $R_l(r_J)$ up to $l = 10$ from latest measurement (8 flybys)
- Lowes' radius: $r_{\text{lowes}} \approx 0.85 r_J$ ($r_J = 6.9894 \times 10^7 \text{m}$)

Questions: with the conductivity profile $\sigma(r)$ varying smoothly,

- meaning of r_{lowes} ? $r_{\text{lowes}} = r_{\text{dyn}}$?
- white source hypothesis valid?
- concept of “dynamo radius” r_{dyn} well-defined?

A numerical model of Jupiter

- spherical shell of radius ratio $r_{\text{in}}/r_{\text{out}} = 0.0963$ (small core)
- rotating fluid with electrical conductivity $\sigma(r)$ driven by buoyancy
- convection forced by secular cooling of the planet
- anelastic: linearise about a hydrostatic adiabatic basic state $(\bar{\rho}, \bar{T}, \bar{p}, \dots)$
- dimensionless numbers: Ra, Pm, Ek, Pr

$$\nabla \cdot (\bar{\rho} \mathbf{u}) = 0$$

$$\frac{Ek}{Pm} \left[\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} \right] + 2\hat{\mathbf{z}} \times \mathbf{u} = -\nabla \Pi' + \frac{1}{\bar{\rho}} (\nabla \times \mathbf{B}) \times \mathbf{B} - \left(\frac{EkRaPm}{Pr} \right) S \frac{d\bar{T}}{dr} \hat{\mathbf{r}} + Ek \frac{\mathbf{F}_\nu}{\bar{\rho}}$$

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{u} \times \mathbf{B}) - \nabla \times (\eta \nabla \times \mathbf{B})$$



$$\bar{\rho} \bar{T} \left(\frac{\partial S}{\partial t} + \mathbf{u} \cdot \nabla S \right) + \frac{Pm}{Pr} \nabla \cdot \mathcal{F}_Q = \frac{Pr}{RaPm} \left(Q_\nu + \frac{1}{Ek} Q_J \right) + \frac{Pm}{Pr} H_S$$

Boundary conditions: no-slip at r_{in} and stress-free at r_{out} , $S(r_{\text{in}}) = 1$ and $S(r_{\text{out}}) = 0$, electrically insulating outside $r_{\text{in}} < r < r_{\text{out}}$. (Jones 2014)

A numerical model of Jupiter

- spherical shell of radius ratio $r_{\text{in}}/r_{\text{out}} = 0.0963$ (small core)
- **rotating** fluid with **electrical conductivity** $\sigma(r)$ driven by **buoyancy**
- convection forced by **secular cooling** of the planet
- **anelastic**: linearise about a hydrostatic adiabatic basic state $(\bar{\rho}, \bar{T}, \bar{p}, \dots)$
- dimensionless numbers: Ra, Pm, Ek, Pr
- **a Jupiter basic state:**


C.A. Jones/Icarus 241 (2014) 148–159




$$Ra = 2 \times 10^7, \quad Ek = 1.5 \times 10^{-5}, \quad Pm = 10, \quad Pr = 0.1$$

radial magnetic field, $B_r(r, \theta, \phi)$


$r = r_{\text{out}}$
dipolar



$r = 0.75r_{\text{out}}$
small scales



Where does the current start flowing?



- average **current** over a spherical surface of radius r

$$\mu_0 \mathbf{j} = \nabla \times \mathbf{B}$$

$$j_{\text{rms}}^2(r, t) \equiv \frac{1}{4\pi} \int_0^{2\pi} \int_0^\pi |\mathbf{j}|^2 \sin \theta \, d\theta \, d\phi$$

- j_{rms} drops quickly but smoothly in the transition region, not clear how to define a characteristic “dynamo radius”

Magnetic energy spectrum, $F_l(r)$

- average magnetic energy over a spherical surface:

$$E_B(r) = \frac{1}{2\mu_0} \frac{1}{4\pi} \oint |\mathbf{B}(r, \theta, \phi)|^2 \sin \theta \, d\theta \, d\phi$$

- Lowes spectrum: recall that if $\mathbf{j} = \mathbf{0}$, we solve $\nabla^2 \Psi = 0$, then


$$2\mu_0 E_B(r) = \sum_{l=1}^{\infty} \left[\left(\frac{a}{r} \right)^{2l+4} (l+1) \sum_{m=0}^l (g_{lm}^2 + h_{lm}^2) \right] = \sum_{l=1}^{\infty} R_l(r)$$

- generally, for the numerical model, $\mathbf{B} \sim \sum_{lm} b_{lm}(r) Y_{lm}(\theta, \phi)$,

$$2\mu_0 E_B(r) = \frac{1}{4\pi} \oint |\mathbf{B}(r, \theta, \phi)|^2 \sin \theta \, d\theta \, d\phi = \sum_{l=1}^{\infty} F_l(r)$$

$$\mathbf{j}(r, \theta, \phi) = \mathbf{0} \text{ exactly} \implies R_l(r) = F_l(r)$$

Magnetic energy spectrum at different depth r




$F_l(r)$: solid lines

$R_l(r)$: circles

- $r > 0.9r_J$: slope of $F_l(r)$ decreases rapidly with r
 $r < 0.9r_J$: $F_l(r)$ maintains the same shape and slope
 \Rightarrow a shift in the dynamics of the system
- $r > 0.9r_J$: $F_l(r) \approx R_l(r)$
 $r < 0.9r_J$: $F_l(r)$ deviates from $R_l(r)$
 \Rightarrow electric current becomes important below $0.9r_J$
- suggests a dynamo radius $r_{\text{dyn}} \approx 0.9r_J$

Spectral slope of $F_l(r)$ and $R_l(r)$



$$\log_{10} F_l(r) \sim -\alpha(r)l$$

$$\log_{10} R_l(r) \sim -\beta(r)l$$


$$R_l(r) = \left(\frac{r_{\text{out}}}{r}\right)^{2l+4} R_l(r_{\text{out}})$$

$$\beta(r) = \beta(r_{\text{out}}) - 2 \log_{10} \frac{r_{\text{out}}}{r}$$

- sharp transition in $\alpha(r)$ indicates $r_{\text{dyn}} = 0.907r_J$
- $F_l(r)$ inside dynamo region is not exactly flat ($\alpha_{\text{dyn}} = 0.024$):
white source assumption is only approximate
- r_{lowes} provides a lower bound to r_{dyn} : $\beta = 0$ at $r_{\text{lowes}} = 0.883$

General picture: $\alpha(r_{\text{out}})$ and α_{dyn} control r_{dyn} and r_{lowes}

Comparison with Juno data



- noise in Juno data \Rightarrow results depend on fitting range
- larger Pm gives smaller α_{dyn} , however $\alpha(r_{\text{out}})$ also becomes smaller $\Rightarrow r_{\text{dyn}}$ remains roughly the same
- $R_l(r_J)$ is shallower in the numerical model than from Juno observation
 - the metallic hydrogen layer could be deeper than predicted by theoretical calculation
 - the existence of a stably stratified layer below the molecular layer
 - our numerical model has more small-scale forcing than Jupiter does

Variation of selected spectral modes with depth

